

**HDAC Inhibitors for Purging HIV-1 from the Latent Reservoir**

Shai Matalon<sup>1</sup>, Thomas A. Rasmussen<sup>2</sup> and Charles A. Dinarello<sup>1</sup>

<sup>1</sup>Department of Medicine, Division of Infectious Diseases, University of Colorado Denver, Aurora, CO 80045 and <sup>2</sup>Department of Infectious Diseases, Aarhus University Hospital, Skejby 8200 Denmark

Address correspondence to Charles A. Dinarello, MD, University of Colorado Denver, Division of Infectious Diseases, 12700 East 19th Ave., B168

Aurora, CO 80045

TEL (303) 724-6172

FAX (303) 724 6178

e-mail [cdinare333@aol.com](mailto:cdinare333@aol.com)

UNCORRECTED PROOF

**Abstract**

A reservoir of latently infected memory CD4<sup>+</sup> T-cells is believed to be the source of HIV-1 re-emergence after discontinuation of anti-retroviral therapy. HIV-1 eradication may depend upon depletion of this reservoir. Integrated HIV-1 is inaccessible for expression, in part due to histone deacetylases. One approach is to exploit the ability of histone deacetylase inhibitors (HDACi) to induce HIV-1 expression from integrated virus. With effective anti-retroviral therapy, newly expressed HIV-1 is incapable of re-infecting naïve cells. With HIV-1 expression, one assumes the infected cell dies and there is a progressive reduction in the size of the reservoir. The concept was tested using the HDACi valproic acid. However valproic acid is weak in inducing HIV-1 from latency in vitro. As such, clinical trials revealed a small or no effect on reducing the number of latently infected T-cells in the peripheral blood. However, new HDAC inhibitors vorinostat, belinostat and givinostat are more effective at targeting specific HDAC's for HIV-1 expression compared to valproic acid. Here, we review studies on HDACi-induced expression of latent HIV-1 with emphasis on new and specific HDAC inhibitors. With increased potency for HIV-1 expression as well as safety and ease of oral administration, new HDACi offer a unique opportunity to deplete the latent reservoir. An additional benefit is the anti-inflammatory properties of HDACi, including down regulation of HIV-1 co-receptor expression.

UNCORRECTED PROOF

**The latent pool of HIV-1 infected cells.**

The source of latent HIV-1 infection is a long-lived pool, most likely the latently infected memory T-cell reservoir, harboring integrated HIV-1 proviral DNA (1-4). (2). Well established techniques are available to quantify the latent pool and it is estimated that one in a million memory T-cells of an HIV-1 positive patient bears a replication competent integrated provirus (5, 6). The latent reservoir of HIV-1 within resting CD4<sup>+</sup> T-cells is established early after the acute infection and the initiation of HAART during this period does not prevent its establishment (6-8). It is an extremely stable reservoir, having a half-life of 6-44 months even in treated patients who are continuously aviremic for long periods of time (6, 9-12). With this prolonged half-life, a complete decay of the reservoir is not expected before 70 years of HAART treatment making eradication improbable. These time frames might be somewhat shorter by starting HAART early during acute infection or by intensifying HAART but are not sufficient as a practical method for eradication (13, 14).

**Early studies on purging HIV-1 from latently infected cells.**

With the development of effective anti-retroviral agents to prevent HIV-1 replication in vivo came the concept that eliminating the reservoir should be achievable without spreading the infection to new uninfected cells. Following viral expression, latently infected cells would then be eliminated by two possible mechanisms. First, it is possible that upon HIV-1 expression, a cytopathic death of the cell would follow. Second, it is also possible that residual HIV-1 specific cytotoxic T-cells would kill cells expressing HIV-1 antigens. The strategy of purging the latent pool of HIV-1 infected CD4<sup>+</sup> T-cells was initially performed using IL-2 and other activators of T-cells such as anti-CD3 antibodies (OKT3). However, although such agents caused a marked activation of the T-cells, there were unacceptable toxicities and also an irreversible decrease in the amount of CD4<sup>+</sup> T-cells (15, 16). Other studies using IL-2 resulted did not reduce the pool (17, 18), perhaps due to a paucity of cell surface receptors for IL-2 on the resting CD4<sup>+</sup> T-cells.

In addition, integrated provirus residing within resting regulatory CD4<sup>+</sup> T-cells (T<sub>regs</sub>) may also have contributed to the lack of effect. The presence of an HIV-1 reservoir within resting T<sub>regs</sub> in patients on prolonged HAART has been detected; in fact, HIV-1 DNA harbouring cells appeared more abundant in this subset than in non-T<sub>regs</sub>, but with a comparable estimated half-life (19). As T<sub>regs</sub> display signs of hyporesponsiveness to cell activation (19), and may inhibit IL-2 production (20), this cellular subset could have added to the viral rebound despite IL-2 treatment (21). However, the anti-inflammatory and immunosuppressive effect of some HDACi, such as SAHA and ITF2357 (22), is in part owing to enhancing the production and suppressive function of T<sub>regs</sub> by promoting FOXP3 acetylation (23). Therefore, it is conceivable, but yet unproven, that potent HDACi treatment may also target the latent viral reservoir within resting T<sub>regs</sub>.

#### **Role of histone acetylation in HIV-1 expression.**

An ideal strategy for viral purging would induce HIV-1 expression without inducing a global T-cell activation. Therefore, small molecules that can directly gain access to DNA and facilitate viral gene expression can circumvent the limitations of cell surface receptor activation. Clearly, HDACi meet these criteria. In general, HDACi are safe, orally active and used widely in medicine. In HIV-1 latency, maintaining histones in their deacetylated form by HDAC's results in densely packed chromatin in the nucleosome with gene expression quiescent. On the other hand, hyperacetylation of specific lysine residues within nucleosomal histones disrupts chromatin binding and allows transcriptional activation of genes. Thus, adding HDACi to cell lines with integrated HIV-1 resulted in expression of HIV-1 without any co-stimulation of cytokines. Table 1 lists some of the reports on HDACi in vitro HIV-1 from cell lines with integrated HIV-1. The human macrophage cell line U1, which is derived from the U937 line served as the model for latent virus in macrophages whereas Ach2 cells were used as the model for integrated HIV-1 in T-cells.

**Which HDAC for HIV-1 expression?**

HDACs are divided into 3 major classes, based on their homolog in yeast. Class I include HDAC-1, 2, 3 and 8. Class II is comprised of HDAC-4, 5, 6, 7, 9 and 10. HDAC-11 has properties of both class I and class II. As shown in Figure 1, the HIV-1 5' long terminal repeat (LTR), containing the promoter and enhancer elements, has binding sites for several transcription factors and is arranged in two nucleosomes (nuc-0 and nuc-1) (24). These nucleosomes are positioned in precise locations with respect to different regulatory elements. During latency, nuc-1 prevents pro-viral transcription, as it remains in the hypoacetylated state. Chromatin immunoprecipitation assays demonstrated that HDAC-1, 2 and 3 have a role in HIV-1 transcriptional regulation (25, 26). Thus, Class I HDACs are recruited to the proviral DNA, in proximity to nuc-1, through the action of DNA binding proteins. NFκB p50 homodimer, AP-4, RBF2, as well as c-Myc and SP-1 recruit Class I HDACs to the LTR, which in turn result in deacetylation of local histones, compaction of the chromatin, and prevention of RNA polymerase-II binding (25, 27-31). In vitro studies revealed that following activation by different stimuli, histone acetyltransferases (HATS) are recruited to the promoter region, leading to acetylation of both H3 and H4 histones. This nuc-1 disruption eventually allows viral transcriptional activation to occur (25, 32, 33).

**Trials with valproic acid to reduce the reservoir of latent virus**

As shown in Table 1, various HDAC inhibitors have been studied in vitro for HIV-1 expression. However, valproic acid (VPA), a carboxylate HDACi which is prescribed for seizures and psychiatric disorders, is the only HDACi which has been combined with HAART in clinical trials. VPA has been administered to HIV-1 infected subjects at doses used to obtain therapeutic blood levels for seizure control while the subjects were maintained throughout the study on the HAART regimen. In the first study by Lehrman et al, enfuvirtide was added to the HAART regimen of 4 HIV-1 positive patients (34). After 4-6 weeks on intensified HAART, VPA was added to the daily regimen for 16 weeks. Plasma levels of VPA were tested and adjusted for 40-100 µg/ml (comparable to 0.25-0.6mM). The frequency of resting cell infection was measured

before and after the augmented therapy and showed a significant reduction in 3 out of 4 patients. However, no measurements were performed between the addition of efavirtide to the regimen and VPA. Therefore, an isolated effect of VPA on reducing the viral latency pool was not established.

In light of that first study, Siliciano et al measured the size of the HIV-1 latent reservoir in 9 patients receiving both HAART and VPA for at least 3 months (35). Levels of latently infected cells isolated from peripheral cells were similar to those seen in patients receiving HAART alone. To exclude the possibility that the patients in the study had an exceptionally large latent pool, the measurement in 7 patients at least 5 months after the initial sample was performed but there was no reduction of the latent reservoir. In fact, in 4 patients the frequencies of HIV-1 gene expression increased. In a French single center study, similar frequencies of CD4<sup>+</sup> T-cells with integrated viral DNA, and CD4<sup>+</sup> T-cells carrying replication-competent virus was observed in both patients who were treated with HAART alone and patients who received HAART plus VPA (36). In 2008, a prospective study compared infection of resting CD4<sup>+</sup> T-cells before and after the addition of VPA to the HAART regimen of 11 HIV-1 positive aviremic patients (37). In only 4 out of 11 patients a depletion of more than 50% in resting CD4<sup>+</sup> T-cells infection was observed, but this effect waned over time, as reported in a follow up study (38). Plasma HIV-1 RNA concentrations were measured also by ultra sensitive real time PCR to a limit of detection of 1 copy/mL (single copy assay). There appeared to be an association between a significant decrease in the number of copies and the absence of low level viremia. Since VPA alone failed to show the desirable effect, the researchers expanded the study and examined the effect of combining intensified anti-viral therapy and the addition of VPA to the regimen of aviremic patients (38). Neither raltegravir nor enfuvirtide, combined with VPA and the baseline HAART, showed a significant decrease.

These studies clearly demonstrate that with or without intensified HAART, the addition of VPA is of limited value in decreasing the reservoir of infected resting CD4<sup>+</sup> T-cells. Nevertheless, the potential of HDAC inhibitors to purge the virus has not been fully exploited. VPA is a weak and non-specific HDACi. VPA does not specifically target HDAC1 or HDAC2, which are thought to be the primary HDAC that

prevent HIV-1 expression. Therefore, the failure to target a specific HDAC in the trials of VPA may account for the lack of an effect. In addition, safety and tolerability are essential requirements for the testing of any experimental drug in patients without a life-threatening disease. For example, we reported that VPA significantly increases HIV-1 expression in the latently infected U1 monocyte/macrophage cell line but at concentrations that would have unacceptable side effects for long-term oral dosing (39).

### **Vorinostat, givinostat, and belinostat**

Using more selective HDACi have been studied in vitro and ex vivo for inducing HIV-1 expression. Suberoylanilide hydroxamic acid (SAHA; generic vorinostat) is a relatively selective inhibitor for Class I HDACs, that is, by inhibiting HDAC 1, 2, 3 and 8 but also with some activity against the Class II HDACs 6, 10 and 11 (39). However, SAHA lacks activity against the Class II HDACs 4, 5, 7 and 8. Although SAHA is approved for the treatment of Cutaneous T-cell Lymphoma, the gastrointestinal side effects and particularly the fall in platelets render the anti-cancer dosing unacceptable for a long-term use in patients with HIV-1(40). Archin et al have examined the ability of SAHA to induce HIV-1 expression in cell lines as well as in virus production from resting CD4<sup>+</sup> T-cells from patients treated with anti-retrovirals (41). In those studies, SAHA induced the in frame marker green fluorescence protein (GFP) mRNA from J89 cells (latently infected Jurkat cell line encoding the enhanced GFP as a marker for Tat-driven HIV LTR expression) at nanomolar concentrations compared to millimolar concentrations for VPA. However, at clinically relevant concentrations of SAHA (340nM) there was no significant increase of GFP mRNA. When SAHA was added to CD4<sup>+</sup> T-cells in vitro at therapeutic concentrations, there was a low level of induction of virus outgrowth in 4 out of 5 patients' cells. Similarly, when used with other types of latently integrated HIV-1 cell lines, the efficacy of SAHA was no different from that of VPA (42).

Similar to SAHA, ITF2357 (generic givinostat) is a hydroxamic acid-containing HDACi that has anti-inflammatory properties in vitro and in vivo as summarized in this issue of Molecular Medicine. At therapeutic plasma levels of 125-250nM, there is no cell toxicity in vitro and only minor thrombocytopenia occurs in patients (43). Givinostat has

been administered for 12 weeks in children with systemic onset juvenile idiopathic arthritis (SoJIA) without side-effects at a therapeutically effective dose of 1.5 mg/kg (44). The use of HDACi in rheumatoid arthritis is reviewed in the present issue of Molecular Medicine (44). We have examined the ability of ITF2357 to induce HIV-1 gene expression in vitro from the latently infected cell lines ACH2 (T-cell line) and U1 (promonocytic line) (39). Givinostat increased p24 by 15-fold in ACH2 cells and, at clinically relevant concentrations, was approximately 10 times more efficient in HIV-1 stimulation than VPA. In U1 cells, VPA failed to double HIV-1 expression as measured by p24 levels, whereas 250nM of ITF2357 induced a 9 fold increase (39).

PXD101 (belinostat), another hydroxamic acid-containing HDACi, has activity against class I and II HDACs at nanomolar concentrations, but slightly less potent than ITF2357 (45). However, as PXD101 has primarily been administered through the intravenous route for treatment of various cancers (46), information on safety, tolerability, and pharmacokinetics using oral administration is still limited (47). Certainly, the side effects from the dosing used in patients with refractory cancer diseases will be unacceptable for HIV-1-infected patients on suppressive HAART. As shown in Figure 2, we compared the ability of VPA, SAHA, ITF2357, and PXD101 to induce HIV-1 expression in the latently infected cell line U1. ITF2357 and PXD101 stimulate HIV-1 expression with a similar potency and both have activity within the low 125-500 nM range. Higher concentrations of SAHA were needed to achieve the same virus production and, as expected, millimolar concentrations of VPA were needed for HIV-1 expression in this system (Fig. 2).

### **The importance of HIV-1 selectivity**

As reviewed above, Class-1 HDACs and specifically HDAC-1, 2 and 3 are considered the major HDACs involved in HIV-1 silencing. Therefore, the use of selective HDAC inhibitors would be a reasonable approach in order to increase its efficacy and at the same time decrease any adverse effects. Specific enzymatic activity assays have been used to compare the selectivity of vorinostat, VPA, and givinostat (39). Analogues of givinostat with different selectivity profile were also examined. Class-1 HDAC selectivity profile correlated directly with HIV-1 stimulation from latently

infected cells U1 and Ach2 cell lines. Whereas VPA showed a weak and non-selective HDAC inhibition, the most selective givinostat analogue showed a 30-fold increase in HIV-1 expression from ACH2 cells. This effect was not due to cell stress. Interestingly, in Ach2 cells, the degree of HDAC-2 inhibition was directly correlated with the induction of HIV-1 gene expression (39). These observations correlate well with the molecular association between the HIV-1 LTR and HDAC-1, 2 and 3.

A recent study by the Margolis group demonstrated that HDACi, which specifically targeted class I HDACs, induced HIV-1 gene expression from cell lines as well as in resting CD4<sup>+</sup> T-cells of aviremic patients (48). An additive effect was seen with inhibition of HDAC-6 as well. These findings support data regarding the selectivity of the class I HDAC-1, 2 and 3 for induction of HIV-1 gene expression. However, preserving the deacetylated status of HDAC-6 prevents infected CD4<sup>+</sup> T-cells HIV-1 from Env-mediated syncytia formation (49). Therefore, analogues that have inhibitory properties targeting HDAC-6 may not be optimal.

### **The effect of HDAC inhibition on HIV-1 co-receptors**

Despite the fact that HDAC inhibitors often increase gene expression, ITF2357 did not increase CCR5 surface expression on CD4<sup>+</sup> T-cells (39). It would be counter-productive if agents stimulating latent virus up-regulated CCR5 and CXCR4 surface expression. In fact, surface expression of CXCR4 on CD4<sup>+</sup> T-cells and CCR5 on monocytes was reduced by 50% by ITF2357 (39). Consistent with decreasing co-receptors surface expression, ITF2357 reduced steady-state mRNA levels of CXCR4 and CCR5 in PBMC, as measured by RT-PCR and by Affimetrix gene expression, respectively. Thus, in a clinical trial of givinostat in HIV-1 infected patients, this property of givinostat may also reduce any re-infection of naïve CD4<sup>+</sup> T-cells.

### **Study design of new HDACi in HIV-1 purging**

Safety is always a consideration when evaluating a drug to treat a disease with no immediate danger to the patient. In testing the hypothesis that HDAC inhibition will purge the latent pool of HIV-1, VPA falls short due to a low level of induced expression

compared to ITF2357. Givostat is safe and effective in humans. In healthy human subjects in a Phase I trial, a single dose of givinostat of 1.5mg/kg resulted in a peak plasma level of 200nM (43). In a Phase II trial in children with active SoJIA, a daily oral dose of givinostat at 1.5mg/kg for 12 weeks exhibited no organ toxicity and achieved significant ( $p<0.01$ ) reduction in parameters of systemic disease as well as the number of painful joints (44). Doses of VPA, on the other hand, when used in order to reach concentrations above 110  $\mu\text{g}/\text{mL}$  in females or 135 $\mu\text{g}/\text{mL}$  in males, carries significant side effects and especially severe thrombocytopenia (50). Thus, higher concentrations of VPA would be impractical to induce more viral expression.

In light of the *in vitro* data suggesting its potency in inducing HIV-1 expression as well as the clinical data suggesting that givinostat is safe, a phase II clinical trial of givinostat would be worthwhile. Administration of givinostat at 1.5 or possibly 1.0 mg/kg in two divided doses may induce a limited number of CD4+ T-cells to express HIV-1. One end-point of such a study is whether after a 12-week course there is a reduction in the inducible pool of latent virus. In addition, it would also be important to determine whether anti-HIV-1 cytotoxic T-cells increased after these courses of givinostat. Although some patients may be reluctant to stop anti-retroviral therapy after courses of HDACi, the *ex vivo* testing of the reservoir would provide evidence that there has been an objective reduction in the latent pool. Surely, the test of the hypothesis that HDACi can reduce the reservoir comes with cessation of anti-retroviral therapy and the duration of time before circulating mRNA levels reappear. Ideally, there would be no return of viral load. Nevertheless, a significant delay in the return of viral load would provide encouragement for the development of specific targeting of HDAC's that assist in maintaining HIV-1 latency.

At another level, givinostat or similar well-tolerated HDACi may provide an additional benefit to HIV-1 infected patients. There are patients being treated with anti-retroviral therapy with less than 50 copies of HIV-1 mRNA but who succumb to all-cause mortality, particularly cardiovascular and renal events (51). In these patients, biomarkers of inflammation, CRP, IL-6 and D-dimer are elevated. The relative risk for death associated with elevated CRP in these patients was 2.0: for elevated IL-6 it was 8.3 and for D-dimer the increased risk was 12.4 (51). In patients who delay anti-retroviral

therapy, there was an expected rise in CD4<sup>+</sup> T-cells upon initiation of anti-retroviral therapy. Although there was also a decrease in the level of D-dimer, there was no change in elevated IL-6 or CRP (52). It is presently unclear whether the decrease in D-dimer will reduce clinical disease; however, the continued elevated IL-6 and CRP levels indicate a continued inflammatory state even in patients being successfully treated with anti-retroviral therapy and who sustain an increased risk for non-HIV-1-associated death. Given the anti-inflammatory properties of HDACi, the markers of systemic inflammation may decrease in a study of HIV-1 purging. On the other hand, it is possible that markers of inflammation may increase as a result of HIV-1 expression and inadequate prevention of re-infection by anti-retroviral therapy.

### Conclusions

Purging the latent reservoir of HIV-1 in resting CD4<sup>+</sup>-T cells by HDAC inhibitors is under active clinical investigation. Use of well-tolerated and selective HDAC inhibitors for HIV-1 purging should be tested in vitro and provide a rationale for clinical studies in aviremic HIV-1 patients on anti-retroviral therapy.

**Acknowledgements.** The authors thank Gregory Pott, Carrie Sailer, Leland Shapiro and Marcel F. Nold for their contributions. Supported by NIH Grant AI-15614 (CAD) and Italfarmaco, SpA, Cinisello Balsamo, Italy.

UNCORRECTED PROOF

## References

1. Imamichi H, *et al.* (2001) Human immunodeficiency virus type 1 quasi species that rebound after discontinuation of highly active antiretroviral therapy are similar to the viral quasi species present before initiation of therapy. *J Infect Dis* 183: 36-50.
2. Zhang L, *et al.* (2000) Genetic characterization of rebounding HIV-1 after cessation of highly active antiretroviral therapy. *J Clin Invest* 106: 839-845.
3. Davey RT, Jr., *et al.* (1999) HIV-1 and T cell dynamics after interruption of highly active antiretroviral therapy (HAART) in patients with a history of sustained viral suppression. *Proc Natl Acad Sci U S A* 96: 15109-15114.
4. Chun TW, *et al.* (2000) Relationship between pre-existing viral reservoirs and the re-emergence of plasma viremia after discontinuation of highly active anti-retroviral therapy. *Nat Med* 6: 757-761.
5. Chun TW, *et al.* (1997) Quantification of latent tissue reservoirs and total body viral load in HIV-1 infection. *Nature* 387: 183-188.
6. Finzi D, *et al.* (1997) Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. *Science* 278: 1295-1300.
7. Chun TW, *et al.* (1998) Early establishment of a pool of latently infected, resting CD4(+) T cells during primary HIV-1 infection. *Proc Natl Acad Sci U S A* 95: 8869-8873.
8. Lori F, *et al.* (1999) Treatment of human immunodeficiency virus infection with hydroxyurea, didanosine, and a protease inhibitor before seroconversion is associated with normalized immune parameters and limited viral reservoir. *J Infect Dis* 180: 1827-1832.
9. Siliciano JD, *et al.* (2003) Long-term follow-up studies confirm the stability of the latent reservoir for HIV-1 in resting CD4+ T cells. *Nat Med* 9: 727-728.
10. Zhang L, *et al.* (1999) Quantifying residual HIV-1 replication in patients receiving combination antiretroviral therapy. *N Engl J Med* 340: 1605-1613.
11. Ramratnam B, *et al.* (2000) The decay of the latent reservoir of replication-competent HIV-1 is inversely correlated with the extent of residual viral replication during prolonged anti-retroviral therapy. *Nat Med* 6: 82-85.
12. Chun TW, *et al.* (2005) HIV-infected individuals receiving effective antiviral therapy for extended periods of time continually replenish their viral reservoir. *J Clin Invest* 115: 3250-3255.
13. Ramratnam B, *et al.* (2004) Intensification of antiretroviral therapy accelerates the decay of the HIV-1 latent reservoir and decreases, but does not eliminate, ongoing virus replication. *J Acquir Immune Defic Syndr* 35: 33-37.
14. Chun TW, *et al.* (2007) Decay of the HIV reservoir in patients receiving antiretroviral therapy for extended periods: implications for eradication of virus. *J Infect Dis* 195: 1762-1764.
15. van Praag RM, *et al.* (2001) OKT3 and IL-2 treatment for purging of the latent HIV-1 reservoir in vivo results in selective long-lasting CD4+ T cell depletion. *J Clin Immunol* 21: 218-226.

16. Prins JM, *et al.* (1999) Immuno-activation with anti-CD3 and recombinant human IL-2 in HIV-1-infected patients on potent antiretroviral therapy. *Aids* 13: 2405-2410.
17. Dybul M, *et al.* (2002) Pilot study of the effects of intermittent interleukin-2 on human immunodeficiency virus (HIV)-specific immune responses in patients treated during recently acquired HIV infection. *J Infect Dis* 185: 61-68.
18. Bowman MC, Archin NM, Margolis DM. (2009) Pharmaceutical approaches to eradication of persistent HIV infection. *Expert Rev Mol Med* 11: e6.
19. Tran TA, *et al.* (2008) Resting regulatory CD4 T cells: a site of HIV persistence in patients on long-term effective antiretroviral therapy. *PLoS One* 3: e3305.
20. Thornton AM, Shevach EM. (1998) CD4+CD25+ immunoregulatory T cells suppress polyclonal T cell activation in vitro by inhibiting interleukin 2 production. *J Exp Med* 188: 287-296.
21. Stellbrink HJ, *et al.* (2002) Effects of interleukin-2 plus highly active antiretroviral therapy on HIV-1 replication and proviral DNA (COSMIC trial). *Aids* 16: 1479-1487.
22. Reddy P, *et al.* (2008) Histone deacetylase inhibition modulates indoleamine 2,3-dioxygenase-dependent DC functions and regulates experimental graft-versus-host disease in mice. *J Clin Invest* 118: 2562-2573.
23. Wang L, de Zoeten EF, Greene MI, Hancock WW. (2009) Immunomodulatory effects of deacetylase inhibitors: therapeutic targeting of FOXP3+ regulatory T cells. *Nat Rev Drug Discov* 8: 969-981.
24. Verdin E, Paras P, Jr., Van Lint C. (1993) Chromatin disruption in the promoter of human immunodeficiency virus type 1 during transcriptional activation. *EMBO J* 12: 3249-3259.
25. Williams SA, *et al.* (2006) NF-kappaB p50 promotes HIV latency through HDAC recruitment and repression of transcriptional initiation. *EMBO J* 25: 139-149.
26. Keedy KS, *et al.* (2009) A limited group of class I histone deacetylases acts to repress human immunodeficiency virus type 1 expression. *J Virol* 83: 4749-4756.
27. Coull JJ, *et al.* (2000) The human factors YY1 and LSF repress the human immunodeficiency virus type 1 long terminal repeat via recruitment of histone deacetylase 1. *J Virol* 74: 6790-6799.
28. Hsia SC, Shi YB. (2002) Chromatin disruption and histone acetylation in regulation of the human immunodeficiency virus type 1 long terminal repeat by thyroid hormone receptor. *Mol Cell Biol* 22: 4043-4052.
29. Jiang G, Espeseth A, Hazuda DJ, Margolis DM. (2007) c-Myc and Sp1 contribute to proviral latency by recruiting histone deacetylase 1 to the human immunodeficiency virus type 1 promoter. *J Virol* 81: 10914-10923.
30. Tyagi M, Karn J. (2007) CBF-1 promotes transcriptional silencing during the establishment of HIV-1 latency. *EMBO J* 26: 4985-4995.
31. Imai K, Okamoto T. (2006) Transcriptional repression of human immunodeficiency virus type 1 by AP-4. *J Biol Chem* 281: 12495-12505.
32. Lusic M, Marcello A, Cereseto A, Giacca M. (2003) Regulation of HIV-1 gene expression by histone acetylation and factor recruitment at the LTR promoter. *EMBO J* 22: 6550-6561.

33. Thierry S, *et al.* (2004) Cell cycle arrest in G2 induces human immunodeficiency virus type 1 transcriptional activation through histone acetylation and recruitment of CBP, NF-kappaB, and c-Jun to the long terminal repeat promoter. *J Virol* 78: 12198-12206.
34. Lehrman G, *et al.* (2005) Depletion of latent HIV-1 infection in vivo: a proof-of-concept study. *Lancet* 366: 549-555.
35. Siliciano JD, *et al.* (2007) Stability of the latent reservoir for HIV-1 in patients receiving valproic acid. *J Infect Dis* 195: 833-836.
36. Sagot-Lerolle N, *et al.* (2008) Prolonged valproic acid treatment does not reduce the size of latent HIV reservoir. *Aids* 22: 1125-1129.
37. Archin NM, *et al.* (2008) Valproic acid without intensified antiviral therapy has limited impact on persistent HIV infection of resting CD4+ T cells. *Aids* 22: 1131-1135.
38. Archin NM, *et al.* (2010) Antiretroviral intensification and valproic acid lack sustained effect on residual HIV-1 viremia or resting CD4+ cell infection. *PLoS One* 5: e9390.
39. Matalon S, *et al.* (2010) The histone deacetylase inhibitor ITF2357 decreases surface CXCR4 and CCR5 expression on CD4(+) T-cells and monocytes and is superior to valproic acid for latent HIV-1 expression in vitro. *J Acquir Immune Defic Syndr* 54: 1-9.
40. Mann BS, Johnson JR, Cohen MH, Justice R, Pazdur R. (2007) FDA approval summary: vorinostat for treatment of advanced primary cutaneous T-cell lymphoma. *Oncologist* 12: 1247-1252.
41. Archin NM, *et al.* (2009) Expression of latent HIV induced by the potent HDAC inhibitor suberoylanilide hydroxamic acid. *AIDS Res Hum Retroviruses* 25: 207-212.
42. Edelstein LC, Micheva-Viteva S, Phelan BD, Dougherty JP. (2009) Short communication: activation of latent HIV type 1 gene expression by suberoylanilide hydroxamic acid (SAHA), an HDAC inhibitor approved for use to treat cutaneous T cell lymphoma. *AIDS Res Hum Retroviruses* 25: 883-887.
43. Furlan A, *et al.* (2011) Pharmacokinetics, safety and inducible cytokine responses during a Phase 1 trial of the histone deacetylase inhibitor ITF2357 (givinostat). *Mol Med* in press.
44. Vojinovic J, Damjanov N. (2011) HDAC inhibition in rheumatoid arthritis and juvenile idiopathic arthritis. *Mol Med* in press.
45. Khan N, *et al.* (2008) Determination of the class and isoform selectivity of small-molecule histone deacetylase inhibitors. *Biochem J* 409: 581-589.
46. Gimsing P, *et al.* (2008) A phase I clinical trial of the histone deacetylase inhibitor belinostat in patients with advanced hematological neoplasia. *Eur J Haematol* 81: 170-176.
47. Mackay HJ, *et al.* Phase II trial of the histone deacetylase inhibitor belinostat in women with platinum resistant epithelial ovarian cancer and micropapillary (LMP) ovarian tumours. *Eur J Cancer* 46: 1573-1579.
48. Archin NM, *et al.* (2009) Expression of latent human immunodeficiency type 1 is induced by novel and selective histone deacetylase inhibitors. *Aids* 23: 1799-1806.

49. Valenzuela-Fernandez A, *et al.* (2005) Histone deacetylase 6 regulates human immunodeficiency virus type 1 infection. *Mol Biol Cell* 16: 5445-5454.
50. Morselli PL, Franco-Morselli R. (1980) Clinical pharmacokinetics of antiepileptic drugs in adults. *Pharmacol Ther* 10: 65-101.
51. Kuller LH, *et al.* (2008) Inflammatory and coagulation biomarkers and mortality in patients with HIV infection. *PLoS Med* 5: e203.
52. Baker JV, *et al.* (2011) Changes in inflammatory and coagulation biomarkers: a randomized comparison of immediate versus deferred antiretroviral therapy in patients with HIV infection. *J Acquir Immune Defic Syndr* 56: 36-43.
53. Laughlin MA, *et al.* (1993) Sodium butyrate treatment of cells latently infected with HIV-1 results in the expression of unspliced viral RNA. *Virology* 196: 496-505.
54. Van Lint C, Emiliani S, Ott M, Verdin E. (1996) Transcriptional activation and chromatin remodeling of the HIV-1 promoter in response to histone acetylation. *EMBO J* 15: 1112-1120.
55. Witvrouw M, *et al.* (1997) Cell type-dependent effect of sodium valproate on human immunodeficiency virus type 1 replication in vitro. *AIDS Res Hum Retroviruses* 13: 187-192.
56. El Kharroubi A, Piras G, Zensen R, Martin MA. (1998) Transcriptional activation of the integrated chromatin-associated human immunodeficiency virus type 1 promoter. *Mol Cell Biol* 18: 2535-2544.
57. Ylisastigui L, Archin NM, Lehrman G, Bosch RJ, Margolis DM. (2004) Coaxing HIV-1 from resting CD4 T cells: histone deacetylase inhibition allows latent viral expression. *Aids* 18: 1101-1108.
58. Contreras X, *et al.* (2009) Suberoylanilide hydroxamic acid reactivates HIV from latently infected cells. *J Biol Chem* 284: 6782-6789.
59. Savarino A, *et al.* (2009) "Shock and kill" effects of class I-selective histone deacetylase inhibitors in combination with the glutathione synthesis inhibitor buthionine sulfoximine in cell line models for HIV-1 quiescence. *Retrovirology* 6: 52.
60. Reuse S, *et al.* (2009) Synergistic activation of HIV-1 expression by deacetylase inhibitors and prostratin: implications for treatment of latent infection. *PLoS One* 4: e6093.
61. Shehu-Xhilaga M, *et al.* (2009) The novel histone deacetylase inhibitors metaccept-1 and metaccept-3 potentially increase HIV-1 transcription in latently infected cells. *Aids* 23: 2047-2050.
62. Choi BS, *et al.* Novel histone deacetylase inhibitors CG05 and CG06 effectively reactivate latently infected HIV-1. *Aids* 24: 609-611.
63. Yin H, Zhang Y, Zhou X, Zhu H. (2010) Histone deacetylase inhibitor Oxamflatin increase HIV-1 transcription by inducing histone modification in latently infected cells. *Mol Biol Rep.*
64. Colin L, Van Lint C. (2009) Molecular control of HIV-1 postintegration latency: implications for the development of new therapeutic strategies. *Retrovirology* 6: 111.

Table 1. Different HDAC inhibitors that have been studied in vitro and ex-vivo for HIV-1 purging <sup>1</sup>.

<u>HDAC Inhibitor</u>	<u>Assay</u>	<u>References</u>
sodium butyrate	TE671/RD cells	(53)
TSA, TPX	U1, ACH2, J49, OM10.1	(54)
VPA	U1, ACH2, HUT-78/IIIB/LAI	(55)
TSA	HeLa	(56)
VPA	RCT	(57)
SAHA	RCT	(58)
SAHA	J89, RCT	(41)
SAHA, IHI	U1, ACH2	(59)
SAHA, VPA	U1, J-Lat	(60)
oxamflatin, VPA SAHA, TSA	ACH2, 8E5, J-Lat	(61)
SAHA, VPA	24STNLESG, Hela	(42)
CG05, CG06	ACH2	(62)
ITF2357, VPA, IHI	ACH2, U1	(39)

TSA, oxamflatin

A7

(63)

<sup>1</sup>TPX, trapoxin; TSA, trichostatin A; VPA, valproic acid; RCT, resting CD4<sup>+</sup> T-cells of aviremic HIV-infected donors; SAHA, suberoylanilide hydroxamic acid; IHI, Investigational HDAC Inhibitors

UNCORRECTED PROOF

## Figure legends

Figure 1. HDAC suppression of HIV-1 expression in latent CD4<sup>+</sup> T-cells. A. In the latent state, HDAC-1, 2 and 3 are recruited to the proviral LTR by different DNA binding proteins. Lysine residues on histones on nuc-1 are hypoacetylated and viral transcription blocked. B. Upon activation of the cell by exogenous stimuli, HDACs are replaced by histone acetyltransferases (HATs), Nuc-1 is hyperacetylated with chromatin remodeling and viral gene transcription. Adapted from (64).

Figure 2. HIV-1 expression in U1 cells stimulated by ITF2357, PDX101, SAHA, or VPA. Mean  $\pm$ SEM p24 levels of two separate experiments for HDAC inhibitors indicated under the horizontal axis. PMA is used as positive control (omitted from figure). ITF2357 was supplied by Italfarmaco S.p.A., PDX101 and SAHA were purchased through Selleck Chemicals LLC.

UNCORRECTED PROOF



